arXiv:1704.00936v1 [math.AP] 4 Apr 2017

NULL CONTROLLABILITY OF A POPULATION DYNAMICS WITH INTERIOR DEGENERACY

IDRISS BOUTAAYAMOU AND YOUNES ECHARROUDI

Abstract. In this paper, we deal with the null controllability of a population dynamics model with an interior degenerate diffusion. To this end, we proved first a new Carleman estimate for the full adjoint system and afterwards we deduce a suitable observability inequality which will be needed to establish the existence of a control acting on a subset of the space which lead the population to extinction in a finite time.

1. Introduction

Consider the system

$$\begin{aligned} \frac{\partial y}{\partial t}+\frac{\partial y}{\partial a}-(k(x)y_x)_x+\mu(t,a,x)y=\vartheta\chi_\omega \qquad & \text{in } Q, \\ y(t,a,1)=y(t,a,0)=0 \qquad & \text{on } (0,T)\times(0,A), \\ y(0,a,x)=y_0(a,x) \qquad & \text{in } Q_A, \\ y(t,0,x)=\int_0^A \beta(t,a,x)y(t,a,x)da \qquad & \text{in } Q_T, \end{aligned} \tag{1.1}$$

where $Q=(0,T)\times(0,A)\times(0,1)$, $Q_A=(0,A)\times(0,1)$, $Q_T=(0,T)\times(0,1)$ and we will denote $q=(0,T)\times(0,A)\times\omega$, where $\omega=(x_1,x_2)\subset\subset(0,1)$ is the region where the control ϑ is acting. This control corresponds to an external supply or to removal of individuals on the subdomain ω. Since the system (1.1) models the dispersion of gene of a given population, then x represents the gene type and $y(t,a,x)$ is the distribution of individuals of age a at time t and of gene type x. The parameters $\beta(t,a,x)$ and $\mu(t,a,x)$ are respectively the natural fertility and mortality rates, A is the maximal age of life and k is the gene dispersion coefficient. $y_0\in L^2(Q_A)$ is the initial distribution of population. Finally, $\int_0^A\beta(t,a,x)y(t,a,x)da$ is the distribution of the newborns of the population that are of gene type x at time t. As usual, we will suppose that no individual reaches the maximal age A. We note that in the most works concerned with the diffusion population dynamics models, x is viewed as the space variable.

The population dynamics models in their different aspects attracted many authors and were investigated from many sides (see for example [18, 20, 21, 25, 28, 29]). Among

2000 *Mathematics Subject Classification.* 35K65, 92D25, 93B05, 93B07.

Key words and phrases. population dynamics model, interior degeneracy, Carleman estimate, observability inequality, null controllability.

Université Ibn Zohr, Faculté polydisciplinaire de Ouarzazate, e-mail: dsboutaayamou@gmail.com.

Département de Mathématiques, Faculté des Sciences Semlalia, Laboratoire LMDP, UMMISCO (IRD-UPMC), B. P. 2390 Marrakech 40000, Maroc, e-mails: yecharroudi@gmail.com.

those questions, we find the null controllability problem or in general the controllability problems for age and space structured population dynamics models which were studied in a intensive literature basing, in general, on the references interested on the controllability of heat equation (see for instance [12, 13, 14, 15, 16, 19] for a different controllability problems of heat equation). In this context, we can cite the pioneering items of V. Barbu and al. in [7], B. Ainseba and S. Anita in [1, 2, 3, 4]. In [7], the authors proved the null controllability for a population dynamics model without diffusion both in the cases of migration and birth control for $T \geq A$ showing directly an appropriate observability inequality for the associated adjoint system and they concluded that in the case of the migration control, only a classes of age was controlled in contrary with the birth control which allows to steer all population to extinction. In [1, 2, 3, 4], the diffusion was taken into account in a age-space structured model and the null controllability of (1.1) for a classes of age was established in the case where $k = 1$ and for any dimension n by means of a weighted estimates called Carleman estimates and exploiting the results gotten for heat equation in [26]. In [5], B. Ainseba and al. studied a more general case allowing the dispersion coefficient to depend on the variable x and verifies $k(0) = 0$ (i.e, the coefficient of dispersion k degenerates at 0). The authors tried to obtain (1.2) in such a situation with $\beta \in L^\infty$ basing on the work done in [6] for the degenerate heat equation to establish a new Carleman estimate for the full adjoint system (2.5) and afterwards his observability inequality. However, the null controllability property of this paper was showed under the condition $T \geq A$ (as in [7]) and this constitutes a restrictiveness on the "optimality" of the control time T since it means, for example, that for a pest population whose the maximal age A may equal to a many days (may be many months or years) we need much time to bring the population to the zero equilibrium. In the same trend and to overcome the condition $T \geq A$, L. Maniar et al in [17] suggested the fixed point technique in which the birth rate β must be in $C^2(Q)$ specially in the proof of [17, Proposition 4.2]. Such a technique consists briefly to demonstrate in a first time the null controllability for an intermediate system with a fertility function $b \in L^2(Q_T)$ instead of $\int_0^A \beta(t, a, x)y(t, a, x)da$ and to achieve the task via a Leray-Schauder Theorem.
Thereby, the main goal of the current paper is to deal with the null controllability property with a minimum of regularity of β (see (2.4)) and a positive small control time T taking into account that k depends on the gene type and degenerates at a point $x_0 \in \omega$, i.e $k(x_0) = 0$, e.g $k(x) = |x - x_0|^\alpha$. To be more accurate, for a fixed $T \in (0, \delta)$ with $\delta \in (0, A)$ small enough, we investigate the existence of a suitable control $\vartheta \in L^2(q)$ which depends on y_0 and δ and such that the associated solution y of (1.1) satisfies

$$y(T, a, x) = 0, \quad \text{a.e. in } (\delta, A) \times (0, 1). \tag{1.2}$$

If $k(x_0) = 0$ in a point $x_0 \in \omega$, we say that (1.1) is a population dynamics model with interior degeneracy. Genetically speaking, the meaning of $k(x_0) = 0$ is that the gene of type $x_0 \in (0, 1)$ can not be transmitted from the studied population to its offspring. This objective will be attained via the classical procedure following the strategy of [22]. On other words, we will establish an appropriate observability inequality for the full adjoint system of (1.1) which is an outcome of a suitable Carleman estimate. We highlight that such a result can be shown if we replace the homogeneous Dirichlet boundary conditions by the ones of Neumann, i.e $y_x(t, a, 0) = y_x(t, a, 1) = 0, (t, a) \in (0, T) \times (0, A)$ using the same way done in [10]. Another interesting null controllability problem of (1.1) can be elaborate using the work of Fragnelli et al. in [23] arising in the case when the potential

term admits an interior singularity belonging to gene type domain.
The remainder of this paper is organized as follows: in Section 2, we will provide the well-posedness of (1.1) and give the proof of the Carleman estimate of its adjoint system. The Section 3 will be devoted to the observability inequality and hence we get the null controllability result (1.2). The last section will take the form of an appendix where we will bring out a Caccioppoli's inequality which plays an important role to show the desired Carleman estimate.

2. **Well-posedness and Carleman estimate results**

2.1. **Well-posedness result.** For this section and for the sequel, we assume that the dispersion coefficient k verifies

$$\begin{cases} \exists x_0 \in (0,1), k \in C([0,1]) \cap C^1([0,1]\backslash\{x_0\}), \ \ k>0 \text{ in } [0,1]\backslash\{x_0\} \text{ and } k(x_0)=0, \\ \exists \gamma \in [0,1) : (x-x_0)k'(x) \leq \gamma k(x), \ x \in [0,1]\backslash\{x_0\}. \end{cases} \tag{2.3}$$

It is well-known in the literature of degenerate problems that there exist two kinds of degeneracy namely the weakly degenerate and the strong degenerate problems, in our study we will restrict ourselves to the first one and this fact explains the choice of $\gamma \in [0,1)$ which in fact are associated to the Dirichlet boundary conditions (see [22, Hypothesis 1.1]). On the other hand, the last hypothesis on k means in the case of $k(x)=|x-x_0|^\alpha$ that $0 \leq \alpha < 1$.
The investigation of (1.2) needs also the following assumptions on the natural rates β and μ

$$\begin{cases} \mu, \beta \in L^\infty(Q), \quad \beta(t,a,x), \mu(t,a,x) \geq 0, \text{ a.e. in } Q, \\ \beta(.,0,.) \equiv 0 \quad \text{a.e. in } (0,T)\times(0,1). \end{cases} \tag{2.4}$$

The last assumption in (2.4) is natural since the newborns are not fertile. Also, it is worth mentioning to point out that, as in [5] we do not need to require that μ satisfies an hypotheses like $\int_0^A \mu(t-s,A-s,x)ds = +\infty, \quad (t,x) \in [0,T]\times[0,1]$ since it does not play any role on the well-posedness result and the computations concerning the proofs of our controllability result as well. However, we will suppose that no individual can reach the maximal age A as mentioned in the introduction. In the same context, we emphasize that in [17], the $L^\infty-$regularity of β is sufficient to prove the well posedness of the studied model which is exactly our case. To this end, we introduce the following weighted Sobolev spaces:

$$\begin{cases} H_k^1(0,1) := \{u \in L^2(0,1) : u \text{ is abs. cont. in } [0,1] : \sqrt{k}u_x \in L^2(0,1), u(1)=u(0)=0\}, \\ H_k^2(0,1) := \Big\{u \in H_k^1(0,1) \ : \ k(x)u_x \in H^1(0,1)\Big\}, \end{cases}$$

endowed respectively with the norms

$$\begin{cases} \|u\|^2_{H_k^1(0,1)} := \|u\|^2_{L^2(0,1)} + \|\sqrt{k}u_x\|^2_{L^2(0,1)}, \quad u \in H_k^1(0,1), \\ \|u\|^2_{H_k^2} := \|u\|^2_{H_k^1(0,1)} + \|(k(x)u_x)_x\|^2_{L^2(0,1)}, \quad u \in H_k^2(0,1). \end{cases}$$

We recall from [24, Theorem 2.2] that the operator $(P, D(P))$ defined by $Pu := (k(x)u_x)_x,\ u \in D(P) = H_k^2(0,1)$, is closed self-adjoint and negative with dense domain in $L^2(0,1)$. Consequently, from [32, Theorem 5] the operator $\mathcal{A} := -\frac{\partial}{\partial a} + P$ generates a C_0-semigroup on the space $L^2((0,A)\times(0,1))$. Then, the following well-posedness result holds.

Theorem 2.1. *Under the assumptions* (2.3) *and* (2.4) *and for all* $\vartheta \in L^2(Q)$ *and* $y_0 \in L^2(Q_A)$*, the system* (1.1) *admits a unique solution* y*. This solution belongs to* $E := C([0,T], L^2((0,A)\times(0,1))) \cap C([0,A], L^2((0,T)\times(0,1))) \cap L^2((0,T)\times(0,A), H_k^1(0,1))$*. Moreover, the solution of* (1.1) *satisfies the following inequality*

$$\sup_{t\in[0,T]} \|y(t)\|^2_{L^2(Q_A)} + \sup_{a\in[0,A]} \|y(a)\|^2_{L^2(Q_T)} + \int_0^1\int_0^A\int_0^T (\sqrt{k(x)}y_x)^2 dtdadx \leq C\left(\int_q \vartheta^2 + \int_{Q_A} y_0^2 dadx\right).$$

2.2. **Carleman estimates results.** As we said in the introduction, we will show the main key of this paper namely the Carleman type inequality. In general, it is well-known that to prove a controllability result of a studied model through this a priori estimate, we must show this last for the associated adjoint system. In our case, this adjoint system takes the following form

$$\begin{aligned} &\frac{\partial w}{\partial t} + \frac{\partial w}{\partial a} + (k(x)w_x)_x - \mu(t,a,x)w = -\beta(t,a,x)w(t,0,x),\\ &w(t,a,1) = w(t,a,0) = 0,\\ &w(T,a,x) = w_T(a,x),\\ &w(t,A,x) = 0, \end{aligned} \tag{2.5}$$

where $T > 0$ and assume that $w_T \in L^2(Q_A)$. Of course, the assumptions (2.3) and (2.4) on k, μ and β are perpetuated. To attaint our goal, we will introduce the following weight functions

$$\begin{cases} \varphi(t,a,x) := \Theta(t,a)\psi(x),\\ \Theta(t,a) := \dfrac{1}{(t(T-t))^4 a^4},\\ \\ \psi(x) := c_1\left(\int_{x_0}^x \frac{r-x_0}{k(r)}dr - c_2\right). \end{cases} \tag{2.6}$$

For the moment, we will assume that $c_2 > \max\{\frac{(1-x_0)^2}{k(1)(2-\gamma)}, \frac{x_0^2}{k(0)(2-\gamma)}\}$ and $c_1 > 0$. A more precise restriction on c_1 will be given later. On the other hand, using the relation satisfied by c_2 and with the aid of [22, Lemma 2.1] one can prove that $\psi(x) < 0\ \forall x \in [0,1]$. Observe also that $\Theta(a,t) \to +\infty$ as $t \to T^-, 0^+$ and $a \to 0^+$.
To demonstrate our Carleman estimate, we require that k fulfills, besides (2.3) the following hypothesis

$$\begin{cases} \exists\theta \in (0,\gamma] \text{ such that :}\\ x \mapsto \frac{k(x)}{|x-x_0|^\theta} \text{ is nonincreasing on the left of } x = x_0\\ \text{ and nondecreasing on the right of } x = x_0, \end{cases} \tag{2.7}$$

where γ is defined by (2.3). The first Carleman estimate result is the following

Proposition 2.2. *Consider the two following systems with $h \in L^2(Q)$*

$$\begin{aligned} \frac{\partial w}{\partial t} + \frac{\partial w}{\partial a} + (k(x)w_x)_x &= h, \\ w(a,t,1) = w(a,t,0) &= 0, \\ w(T,a,x) &= w_T(a,x), \\ w(t,A,x) &= 0, \end{aligned} \tag{2.8}$$

and

$$\begin{aligned} \frac{\partial w}{\partial t} + \frac{\partial w}{\partial a} + (k(x)w_x)_x - \mu(t,a,x)w &= h, \\ w(t,a,1) = w(t,a,0) &= 0, \\ w(T,a,x) &= w_T(a,x), \\ w(t,A,x) &= 0. \end{aligned} \tag{2.9}$$

Then, there exist two positive constants C and s_0, such that every solutions of (2.8) *and* (2.9) *satisfy, for all $s \geq s_0$, the following inequality*

$$\begin{aligned} &s^3 \int_Q \Theta^3 \frac{(x-x_0)^2}{k(x)} w^2 e^{2s\varphi} dtdadx + s \int_Q \Theta k(x) w_x^2 e^{2s\varphi} dtdadx \\ &\leq C \left(\int_Q |\, h \,|^2 \, e^{2s\varphi} dtdadx + s \int_0^A \int_0^T [k\Theta e^{2s\varphi}(x-x_0) w_x^2]_{x=0}^{x=1} dtda \right). \end{aligned} \tag{2.10}$$

Proof. Firstly, we will prove (2.10) for system (2.8) and replacing h by $h + \mu w$ we will get the same inequality for (2.9). So, let w be the solution of (2.8) and put

$$\nu(t,a,x) := e^{s\varphi(t,a,x)} w(t,a,x)$$

. Then, ν satisfies the following system

$$\begin{aligned} L_s^+ \nu + L_s^- \nu &= e^{s\varphi(t,a,x)} h, \\ \nu(t,a,1) = \nu(t,a,0) &= 0, \\ \nu(T,a,x) = \nu(0,a,x) &= 0, \\ \nu(t,A,x) = \nu(t,0,x) &= 0, \end{aligned} \tag{2.11}$$

where

$$L_s^+ \nu := (k(x)\nu_x)_x - s(\varphi_a + \varphi_t)\nu + s^2 \varphi_x^2 k(x) \nu,$$

and

$$L_s^- \nu := \nu_t + \nu_a - 2sk(x)\varphi_x \nu_x - s(k(x)\varphi_x)_x \nu.$$

Passing to the norm in (2.11), one has

$$\|L_s^+ \nu\|^2_{L^2(Q)} + \|L_s^- \nu\|^2_{L^2(Q)} + 2 < L_s^+ \nu, L_s^- \nu >= \|e^{s\varphi(a,t,x)} h\|^2_{L^2(Q)},$$

where $< ., . >$ denotes here the inner product in $L^2(Q)$. Then, the proof of step one is based on the calculus of the inner product $< L_s^+ \nu, L_s^- \nu >$ whose a first expression is given in the following lemma.

Lemma 2.3. *The following identity holds*

$$< L_s^+\nu, L_s^-\nu >= S_1 + S_2,$$

with

$$\begin{aligned}
S_1 &= s\int_Q (k(x)\nu_x)^2\varphi_{xx}dtdadx - s^3\int_Q (k(x)\varphi_x)_x k(x)\varphi_x^2\nu^2 dtdadx + s^2\int_Q (\varphi_a+\varphi_t)(k(x)\varphi_x)_x\nu^2 dtdadx\\
&+ s\int_Q k(x)\nu_x((k(x)\varphi_x)_{xx}\nu + (k(x)\varphi_x)_x\nu_x)dtdadx + s^3\int_Q (k^2\varphi_x^3)_x\nu^2 dtdadx\\
&- s^2\int_Q (k(x)(\varphi_a+\varphi_t)\varphi_x)_x\nu^2 dtdadx + \frac{s}{2}\int_Q (\varphi_{at}+\varphi_{tt})\nu^2 dtdadx - \frac{s^2}{2}\int_Q (\varphi_x^2)_t k(x)\nu^2 dtdadx\\
&+ \frac{s}{2}\int_Q (\varphi_{at}+\varphi_{aa})\nu^2 dtdadx - \frac{s^2}{2}\int_Q (\varphi_x^2)_a k(x)\nu^2 dtdadx,
\end{aligned}$$

and

$$\begin{aligned}
S_2 &= \int_0^A\int_0^T [k(x)\nu_x\nu_a]_0^1 dtda + \int_0^A\int_0^T [k(x)\nu_x\nu_t]_0^1 dtda\\
&+s^2\int_0^A\int_0^T [k(x)\varphi_x(\varphi_a+\varphi_t)\nu^2]_0^1 dtda - s^3\int_0^A\int_0^T [k^2(x)\varphi_x^3\nu^2]_0^1 dtda\\
&-s\int_0^A\int_0^T [k(x)\nu\nu_x(k(x)\varphi_x)_x]_0^1 dtda - s\int_0^A\int_0^T [(k(x)\nu_x)^2\varphi_x]_0^1 dtda.
\end{aligned}$$

For the proof of Lemma 2.3, see the one of [17, Lemma 3.2]. The previous expressions of S_1 and S_2 can be simplified using of the functions φ and ψ given in (2.6) and also the homogeneous Dirichlet boundary conditions satisfied by ν. Hence, one has

$$\begin{aligned}
S_1 &= \frac{s}{2}\int_Q (\Theta_{aa}+\Theta_{tt})\psi\nu^2 dxdtda + s\int_Q \Theta_{ta}\psi\nu^2 dtdadx\\
&+sc_1\int_Q \Theta(2k(x)-(x-x_0)k'(x))\nu_x^2 dtdadx - 2s^2\int_Q \Theta c_1^2\frac{(x-x_0)^2}{k(x)}(\Theta_a+\Theta_t)\nu^2 dtdadx\\
&+s^3\int_Q \Theta^3 c_1^3\left(\frac{x-x_0}{k(x)})^2(2k(x)-(x-x_0)k'(x)\right)\nu^2 dtdadx, \qquad (2.12)
\end{aligned}$$

and

$$S_2 = -sc_1\int_0^A\int_0^T [k\Theta e^{2s\varphi}(x-x_0)\nu_x^2]_{x=0}^{x=1} dtda.$$

Accordingly,

$$\begin{aligned}
< L_s^+\nu, L_s^-\nu > &= \frac{s}{2}\int_Q (\Theta_{aa}+\Theta_{tt})\psi\nu^2 dxdtda + s\int_Q \Theta_{ta}\psi\nu^2 dtdadx\\
&+sc_1\int_Q \Theta(2k(x)-(x-x_0)k'(x))\nu_x^2 dtdadx - 2s^2\int_Q \Theta c_1^2\frac{(x-x_0)^2}{k(x)}(\Theta_a+\Theta_t)\nu^2 dtdadx\\
&+s^3\int_Q \Theta^3 c_1^3\left(\frac{x-x_0}{k(x)}\right)^2(2k(x)-(x-x_0)k'(x))\nu^2 dtdadx \qquad (2.13)\\
&-sc_1\int_0^A\int_0^T [k\Theta e^{2s\varphi}(x-x_0)w_x^2]_{x=0}^{x=1} dtda.
\end{aligned}$$

Thanks to the third assumption in (2.3), we have

$$S_1 \geq \frac{s}{2}\int_Q (\Theta_{aa}+\Theta_{tt})\psi\nu^2 dtdadx + s\int_Q \Theta_{ta}\psi\nu^2 dtdadx + sc_1\int_Q \Theta k(x)\nu_x^2 dtdadx \quad (2.14)$$
$$-2s^2\int_Q \Theta c_1^2\frac{(x-x_0)^2}{k(x)}(\Theta_a+\Theta_t)\nu^2 dtdadx + s^3\int_Q \Theta^3 c_1^3\frac{(x-x_0)^2}{k(x)}\nu^2 dtdadx.$$

Observe that $|\Theta(\Theta_a+\Theta_t)| \leq c\Theta^3$, to infer, for s quite large that

$$\left|-2s^2\int_Q \Theta c_1^2\frac{(x-x_0)^2}{k(x)}(\Theta_a+\Theta_t)\nu^2 dtdadx\right|$$
$$\leq 2s^2c_1^2c\int_Q \frac{(x-x_0)^2}{k(x)}\Theta^3\nu^2 dtdadx \leq \frac{c_1^3}{4}s^3\int_Q \frac{(x-x_0)^2}{k(x)}\Theta^3\nu^2 dtdadx. \quad (2.15)$$

On the other hand, we have $r \mapsto \frac{|r-x_0|^\gamma}{k(r)}$ is nondecreasing in the right of x_0.
Then,

$$|\psi(x)| = |c_1 l(x) - c_1c_2| \leq c_1\left|\int_{x_0}^x \frac{r-x_0}{k(r)}dr\right| + c_1c_2 \leq c_1c_2 + c_1\frac{(1-x_0)^2}{k(1)(2-\gamma)}$$
$$\leq \frac{c_1}{(2-\gamma)k(1)} + c_1c_2.$$

A simple computations allow us to check that $\Theta_{aa}+\Theta_{tt}+|\Theta_{ta}| \leq C_1\Theta^{\frac{3}{2}}$. This yields

$$\left|\frac{s}{2}\int_Q (\Theta_{aa}+\Theta_{tt})\psi\nu^2 dtdadx + s\int_Q \Theta_{ta}\psi\nu^2 dtdadx\right|$$
$$\leq s\left(\frac{c_1}{(2-\gamma)k(1)} + c_1c_2\right)\int_Q \left(\frac{\Theta_{aa}+\Theta_{tt}}{2} + |\Theta_{ta}|\right)\nu^2 dtdadx$$
$$\leq Ms\left(\frac{c_1}{(2-\gamma)k(1)} + c_1c_2\right)\int_Q \Theta^{\frac{3}{2}}\nu^2 dtdadx. \quad (2.16)$$

It remains now to bound the term $|\int_Q \Theta^{\frac{3}{2}}\nu^2 dtdadx|$. Using the generalized Young inequality we obtain

$$\left|\int_0^1 \Theta^{\frac{3}{2}}\nu^2 dx\right|$$
$$= \left|\int_0^1 \left(\Theta\frac{k^{\frac{1}{3}}}{|x-x_0|^{\frac{2}{3}}}\nu^2\right)^{\frac{3}{4}}\left(\Theta^3\frac{|x-x_0|^2}{k}\nu^2\right)^{\frac{1}{4}}dx\right|$$
$$\leq \frac{3\epsilon}{4}\int_0^1 \Theta\frac{k^{\frac{1}{3}}}{|x-x_0|^{\frac{2}{3}}}\nu^2 dx + \frac{1}{4\epsilon}\int_0^1 \Theta^3\frac{|x-x_0|^2}{k}\nu^2 dx \quad (2.17)$$

Put

$$p(x) = (k(x)|x-x_0|^4)^{\frac{1}{3}}. \quad (2.18)$$

By hypothesis (2.7), one can check that $x \mapsto \frac{p(x)}{|x-x_0|^q}$, with $q := \frac{4+\theta}{3} \in (1,2)$ is nonincreasing on the left of $x = x_0$ and nondecreasing on the right of $x = x_0$. Furthermore, we have

$\frac{k^{\frac{1}{3}}}{|x-x_0|^{\frac{2}{3}}} = \frac{p(x)}{(x-x_0)^2}$ and there exists $C_2 > 0$ such that $p(x) < C_2 k(x)$.
Hence, by mean of Hardy-Poincaré inequality (see [22, Proposition 2.3]), we conclude that

$$\begin{aligned}\int_0^1 \Theta \frac{k^{\frac{1}{3}}}{|x-x_0|^{\frac{2}{3}}}\nu^2 dx &= \int_0^1 \Theta \frac{p(x)}{(x-x_0)^2}\nu^2 dx \\ &\leq C\int_0^1 \Theta p \nu_x^2 dx \\ &\leq CC_2 \int_0^1 \Theta k \nu_x^2 dx, \end{aligned} \tag{2.19}$$

where $C > 0$ is the constant of Hardy-Poincaré. Combining (2.17) and (2.19), we get

$$\left|\int_0^1 \Theta^{\frac{3}{2}}\nu^2 dx\right| \leq \frac{3\epsilon}{4}C_3 \int_Q \Theta k \nu_x^2 dtdadx + \frac{1}{4\epsilon}\int_Q \Theta^3 \frac{|x-x_0|^2}{k}\nu^2 dtdadx. \tag{2.20}$$

Hence, (2.16) and (2.20) lead to

$$\begin{aligned}&\left|\frac{s}{2}\int_Q (\Theta_{aa}+\Theta_{tt})\psi\nu^2 dtdadx + s\int_Q \Theta_{ta}\psi\nu^2 dtdadx\right| \\ &\leq sc_1C_4\epsilon \int_Q \Theta k\nu_x^2 dtdadx + \frac{sc_1C_5}{4\epsilon}\int_Q \Theta^3\frac{|x-x_0|^2}{k}\nu^2 dtdadx. \end{aligned} \tag{2.21}$$

Taking ϵ small enough and s quite large, we conclude that

$$\begin{aligned}&\left|\frac{s}{2}\int_Q (\Theta_{aa}+\Theta_{tt})\psi\nu^2 dtdadx + s\int_Q \Theta_{ta}\psi\nu^2 dtdadx\right| \\ &\leq \frac{sc_1}{4}\int_Q \Theta k\nu_x^2 dtdadx + \frac{s^3c_1^3}{4}\int_Q \Theta^3\frac{|x-x_0|^2}{k}\nu^2 dtdadx. \end{aligned} \tag{2.22}$$

Taking into account the relations (2.14) and (2.15) we arrive to

$$S_1 \geq K_1 s^3 \int_Q \Theta^3 \frac{|x-x_0|^2}{k}\nu^2 dtdadx + K_2 s\int_Q \Theta k\nu_x^2 dtdadx \tag{2.23}$$

Hence,

$$\begin{aligned}&2 < L_s^+\nu, L_s^-\nu > \geq m\left(s^3\int_Q \Theta^3\frac{|x-x_0|^2}{k}\nu^2 dtdadx + s\int_Q \Theta k\nu_x^2 dtdadx\right) \\ &-2sc_1\int_0^A\int_0^T [k\Theta e^{2s\varphi}(x-x_0)\nu_x^2]_{x=0}^{x=1} dtda \end{aligned} \tag{2.24}$$

This steers to the following Carleman estimate verified by ν solution of (2.11)

$$\begin{aligned}&s^3\int_Q \Theta^3\frac{|x-x_0|^2}{k}\nu^2 dtdadx + s\int_Q \Theta k\nu_x^2 dtdadx \\ &\leq C_6\left(\int_Q h^2 e^{2s\varphi} dtdadx + s\int_0^A\int_0^T [k\Theta e^{2s\varphi}(x-x_0)\nu_x^2]_{x=0}^{x=1} dtda\right) \end{aligned} \tag{2.25}$$

By the definition of ν we infer that

$$\nu_x = s\varphi_x e^{s\varphi}w + e^{s\varphi}w_x, \quad e^{2s\varphi}w_x^2 \leq 2(\nu_x^2 + s^2\varphi_x^2\nu^2). \tag{2.26}$$

Finally, the Carleman estimate (2.10) of (2.8) is obtained.
Now, If we apply the same inequality of Hardy-Poincaré in a similar way as before to the function $\nu := e^{s\varphi}w$, taking into account the hypothesis on μ assumed in (2.4), using the Carleman type inequality (2.10) for the function $h + \mu w$ and taking s quite large we achieve the Proposition 2.2. □

With the aid of the estimate (2.10) and Caccioppoli's inequality (4.72), we can now show a ω-local Carleman estimate for the system (2.9). This result will be useful to show our main Carleman estimate replacing the second term h by $-\beta(t,a,x)w(t,0,x)$. To this end, we introduce the following weight functions:

$$\begin{cases} \Phi(t,a,x) := \Theta(t,a)\Psi(x), \\ \Psi(x) = e^{\kappa\sigma(x)} - e^{2\kappa\|\sigma\|_\infty}, \end{cases} \tag{2.27}$$

where Θ is given by (2.6), $\kappa > 0$ and σ is the function given by

$$\begin{cases} \sigma \in C^2([0,1]), \sigma(x) > 0 \text{ in } (0,1), \sigma(0) = \sigma(1) = 0, \\ \sigma_x(x) \neq 0 \text{ in } [0,1]\backslash\omega_0, \end{cases} \tag{2.28}$$

where $\omega_0 \Subset \omega$ is an open subset. The existence of the function σ is proved in [26].
On the other hand by the definition of φ (2.6) and taking

$$c_1 \geq \max\left(\frac{k(1)(2-\gamma)(e^{2\kappa\|\sigma\|_\infty}-1)}{c_2 k(1)(2-\gamma)-(1-x_0)^2}, \frac{k(0)(2-\gamma)(e^{2\kappa\|\sigma\|_\infty}-1)}{c_2 k(0)(2-\gamma)-x_0^2}\right), \tag{2.29}$$

one can prove that

$$\varphi \leq \Phi. \tag{2.30}$$

Our theorem is stated as follows

Theorem 2.4. *Assume that the assumptions* (2.3), (2.4) *and* (2.7) *hold. Let $A > 0$ and $T > 0$ be given. Then, there exist positive constants C and s_0 such that for all $s \geq s_0$, every solution w of* (2.9) *satisfies*

$$\int_Q (s\Theta k w_x^2 + s^3\Theta^3\frac{(x-x_0)^2}{k}w^2)e^{2s\varphi}dtdadx \leq C\left(\int_Q h^2 e^{2s\Phi}dtdadx + \int_q s^3\Theta^3 w^2 e^{2s\Phi}dtdadx\right), \tag{2.31}$$

To prove this theorem, we need the following result which represents the Carleman estimate of nondegenerate population dynamics systems. This inequality is stated as follows

Proposition 2.5. *Let us consider the following system*

$$\begin{aligned} \frac{\partial z}{\partial t} + \frac{\partial z}{\partial a} + (k(x)z_x)_x - c(t,a,x)z = h \qquad & \text{in } Q_b, \\ z(t,a,b_1) = z(t,a,b_2) = 0 \qquad & \text{on } (0,T)\times(0,A), \end{aligned} \tag{2.32}$$

where $Q_b := (0,T)\times(0,A)\times(b_1,b_2)$, $(b_1,b_2) \subset [0,x_0)$, or $(b_1,b_2) \subset (x_0,1]$, $h \in L^2(Q_b)$, $k \in C^1([0,1])$ is a strictly positive function and $c \in L^\infty(Q_b)$. Then, there exist two positive constants C and s_0, such that for any $s \geq s_0$, z verifies the following estimate

$$\int_{Q_b} (s^3\phi^3 z^2 + s\phi z_x^2)e^{2s\Phi}dtdadx$$

$$\leq C\left(\int_{Q_b} h^2 e^{2s\Phi} dtdadx + \int_\omega \int_0^A \int_0^T s^3\phi^3 z^2 e^{2s\Phi} dtdadx\right), \tag{2.33}$$

where

$$\phi(t,a,x) = \Theta(t,a)e^{\kappa\sigma(x)}, \tag{2.34}$$

Θ *and* Φ *are defined by* (2.27) *and* σ *by* (2.28).

Before giving the proof of Theorem 2.4, we note that a similar result was demonstrated in [2, Lemma 2.1] in the case when k is a positive constant, for any dimension n without the source term h and with the weight function $\Theta(t,a) = \frac{1}{t(T-t)a}$. By careful computations, the same proof can be adapted to (2.33) where k is a positive general nondegenerate coefficient, with our weight function $\Theta(t,a) = \frac{1}{t^4(T-t)^4a^4}$ and the source term h.

Proof. Let us introduce the smooth cut-off function $\xi : \mathbb{R} \to \mathbb{R}$ defined by

$$\begin{cases} 0 \leq \xi(x) \leq 1, & x \in [0,1], \\ \xi(x) = 1, & x \in [\lambda_1, \lambda_2], \\ \xi(x) = 0, & x \in [0,1]\backslash\omega, \end{cases} \tag{2.35}$$

where $\lambda_1 = \frac{x_1+2x_0}{3}$ and $\lambda_2 = \frac{x_0+2x_2}{3}$.
Let w be the solution of (2.9) and define $v := \xi w$. Then, v satisfies the following system

$$\begin{aligned} \frac{\partial v}{\partial t} + \frac{\partial v}{\partial a} + (k(x)v_x)_x - \mu(t,a,x)v &= \overline{h}, \\ v(t,a,1) = v(t,a,0) &= 0, \\ v(T,a,x) &= \xi w_T(a,x), \\ v(t,A,x) &= 0, \end{aligned} \tag{2.36}$$

where $\overline{h} := \xi h + (k(x)\xi_x w)_x + k(x)w_x\xi_x$.
Using Carleman estimate (2.10) and the definition of ξ, one has

$$\int_Q (s\Theta k v_x^2 + s^3\Theta^3 \frac{(x-x_0)^2}{k} v^2) e^{2s\varphi} dtdadx \leq C\int_Q \overline{h}^2 e^{2s\varphi} dtdadx. \tag{2.37}$$

On the other hand, using again the definition of ξ we can check readily that

$$\begin{aligned} &\int_{\lambda_1}^{\lambda_2}\int_0^A\int_0^T (s\Theta k v_x^2 + s^3\Theta^3 \frac{(x-x_0)^2}{k} v^2) e^{2s\varphi} dtdadx \\ = &\int_{\lambda_1}^{\lambda_2}\int_0^A\int_0^T (s\Theta k w_x^2 + s^3\Theta^3 \frac{(x-x_0)^2}{k} w^2) e^{2s\varphi} dtdadx. \end{aligned} \tag{2.38}$$

Therefore, combining (2.37) and (2.38) we have

$$\begin{aligned} &\int_{\lambda_1}^{\lambda_2}\int_0^A\int_0^T (s\Theta k w_x^2 + s^3\Theta^3 \frac{(x-x_0)^2}{k} w^2) e^{2s\varphi} dtdadx \\ &\leq C\int_Q \overline{h}^2 e^{2s\varphi} dtdadx. \end{aligned} \tag{2.39}$$

Hence by means of Caccioppoli's inequality (4.72) and (2.39), we conclude that

$$\int_{\lambda_1}^{\lambda_2}\int_0^A\int_0^T (s\Theta k w_x^2 + s^3\Theta^3\frac{(x-x_0)^2}{k}w^2)e^{2s\varphi}dtdadx$$
$$\leq C\left(\int_Q h^2 e^{2s\varphi}dtdadx + \int_q s^2\Theta^2 w^2 e^{2s\varphi}dtdadx\right), \tag{2.40}$$

where $\omega^{'}$ of Lemma 4.1 here is exactly $(x_1,\lambda_1)\cup(\lambda_2,x_2)$.
Now, let $z := \eta w$, with η is the smooth cut-off function defined by

$$\begin{cases} 0\leq \eta(x)\leq 1, & x\in[0,1],\\ \eta(x)=0, & x\in[0,\frac{\lambda_3+2\lambda_2}{3}],\\ \eta(x)=1, & x\in[\lambda_2,1], \end{cases} \tag{2.41}$$

where $\lambda_3 = \frac{x_2+2x_0}{3}$. We can observe easily that $\lambda_3 < \frac{\lambda_3+2\lambda_2}{3} < \lambda_2$. Then, z satisfies the following population dynamics equation

$$\begin{aligned} \frac{\partial z}{\partial t}+\frac{\partial z}{\partial a}+k(x)z_{xx}+k^{'}(x)z_x-\mu(t,a,x)z &= \widetilde{h}, \quad \text{in } (\lambda_3,1) \\ z(t,a,1)=z(t,a,\lambda_3) &= 0, \quad \text{in },(0,T)\times(0,A) \end{aligned} \tag{2.42}$$

where $\widetilde{h} := \eta h + (k(x)\eta_x w)_x + k(x)w_x\eta_x$.
By assumption on k, we have $k(x) > 0,\ \ x\in(\lambda_3,1)$. Hence, (2.42) is a nondegenerate model. In this case, applying Proposition 2.5 to the function $\widetilde{h}$ with $b_1=\lambda_3$ and $b_2=1$ and using again Caccioppoli's inequality (4.72), we infer that

$$\begin{aligned} &\int_{\lambda_3}^1\int_0^A\int_0^T (s^3\phi^3 z^2 + s\phi z_x^2)e^{2s\Phi}dtdadx \\ &\leq C\left(\int_{\lambda_3}^1\int_0^A\int_0^T(\eta h+(k\eta_x w)_x+k\eta_x w_x)^2 e^{2s\Phi}dtdadx + \int_\omega\int_0^A\int_0^T s^3\Theta^3 w^2 e^{2s\Phi}dtdadx\right)\\ &\leq \widetilde{C}\left(\int_Q h^2e^{2s\Phi}+((k\eta_x w)_x+k\eta_x w_x)^2e^{2s\Phi}dtdadx + \int_\omega\int_0^A\int_0^T s^3\Theta^3w^2e^{2s\Phi}dtdadx\right)\\ &\leq \widetilde{C}(\int_Q h^2e^{2s\Phi}dtdadx + \int_{\omega'}\int_0^A\int_0^T (8(k\eta_x)^2w_x^2+2((k\eta_x)_x)^2w^2)e^{2s\Phi}dtdadx\\ &+\int_\omega\int_0^A\int_0^T s^3\Theta^3w^2e^{2s\Phi}dtdadx)\\ &\leq \widetilde{C}_1\left(\int_Q h^2e^{2s\Phi}dtdadx+\int_{\omega'}\int_0^A\int_0^T(w_x^2+w^2)e^{2s\Phi}dtdadx+\int_\omega\int_0^A\int_0^T s^3\Theta^3w^2e^{2s\Phi}dtdadx\right)\\ &\leq \widetilde{C}_2\left(\int_Q h^2e^{2s\Phi}dtdadx+\int_\omega\int_0^A\int_0^T s^3\Theta^3w^2e^{2s\Phi}dtdadx\right), \end{aligned} \tag{2.43}$$

with $\omega^{'} := (\frac{\lambda_3+2\lambda_2}{3},\lambda_2)$. By the restriction (2.30) there exists $c_3>0$ such that, for $(t,a,x)\in[0,T]\times[0,A]\times[\lambda_3,1]$, we have

$$\Theta k(x)e^{2s\varphi}\leq c_3\phi e^{2s\Phi},\quad \Theta^3\frac{(x-x_0)^2}{k(x)}e^{2s\varphi}\leq c_3\phi^3e^{2s\Phi}.$$

Then,

$$\int_{\lambda_3}^{1}\int_0^A\int_0^T (s\Theta k z_x^2 + s^3\Theta^3\frac{(x-x_0)^2}{k}z^2)e^{2s\varphi}dtdadx$$
$$\leq c_3\int_{\lambda_3}^{1}\int_0^A\int_0^T (s^3\phi^3 z^2 + s\phi z_x^2)e^{2s\Phi}dtdadx. \tag{2.44}$$

This inequality together with (2.43) lead to

$$\int_{\lambda_3}^{1}\int_0^A\int_0^T (s\Theta k z_x^2 + s^3\Theta^3\frac{(x-x_0)^2}{k}z^2)e^{2s\varphi}dtdadx$$
$$\leq \widetilde{c_3}\left(\int_Q h^2 e^{2s\Phi}dtdadx + \int_\omega\int_0^A\int_0^T s^3\Theta^3 w^2 e^{2s\Phi}dtdadx\right). \tag{2.45}$$

Taking into account the definition of η (2.41), we can say that

$$\int_{\lambda_2}^{1}\int_0^A\int_0^T (s\Theta k z_x^2 + s^3\Theta^3\frac{(x-x_0)^2}{k}z^2)e^{2s\varphi}dtdadx$$
$$= \int_{\lambda_2}^{1}\int_0^A\int_0^T (s\Theta k w_x^2 + s^3\Theta^3\frac{(x-x_0)^2}{k}w^2)e^{2s\varphi}dtdadx. \tag{2.46}$$

Hence,

$$\int_{\lambda_2}^{1}\int_0^A\int_0^T (s\Theta k w_x^2 + s^3\Theta^3\frac{(x-x_0)^2}{k}w^2)e^{2s\varphi}dtdadx$$
$$\leq \widetilde{c_3}\left(\int_Q h^2 e^{2s\Phi}dtdadx + \int_\omega\int_0^A\int_0^T s^3\Theta^3 w^2 e^{2s\Phi}dtdadx\right), \tag{2.47}$$

as a consequence of (2.45) and (2.46). Arguing in the same way for $(0,\lambda_1)$, one can show that

$$\int_{0}^{\lambda_1}\int_0^A\int_0^T (s\Theta k w_x^2 + s^3\Theta^3\frac{(x-x_0)^2}{k}w^2)e^{2s\varphi}dtdadx$$
$$\leq \widetilde{c_3}\left(\int_Q h^2 e^{2s\Phi}dtdadx + \int_\omega\int_0^A\int_0^T s^3\Theta^3 w^2 e^{2s\Phi}dtdadx\right), \tag{2.48}$$

Finally, summing the inequalities (2.40), (2.47) and (2.48) side by side, taking s quite large and using again the restriction on c_1 (2.30) we arrive to

$$\int_{0}^{1}\int_0^A\int_0^T (s\Theta k w_x^2 + s^3\Theta^3\frac{(x-x_0)^2}{k}w^2)e^{2s\varphi}dtdadx$$
$$\leq \widetilde{c_3}\left(\int_Q h^2 e^{2s\Phi}dtdadx + \int_\omega\int_0^A\int_0^T s^3\Theta^3 w^2 e^{2s\Phi}dtdadx\right), \tag{2.49}$$

and this is exactly the desired estimate (2.31). □

Before to provide the main Carleman estimate, we make the following remarks:

Remark 2.6. *1/ The proof of our distributed-Carleman estimate* (2.31) *is based on the cut-off functions and given by two different weighted functions* φ *and* Φ*, in addition by* (2.30) *there is no positive constant* C *such that:*

$$e^{2s\Phi} \leq Ce^{2s\varphi}.$$

2/ Our proof is not based on the reflection method used for the proof of [22, Lemma 4.1] *which is needed to eliminate the boundary term arising in the classical Carleman estimate for nondegenerate heat equation.*

By the Carleman estimate (2.31), we are able to show the following ω-Carleman estimate for the full adjoint system (2.5)

Theorem 2.7. *Assume that the assumptions* (2.3), (2.4) *and* (2.7) *hold. Let* $A > 0$ *and* $T > 0$ *be given such that* $0 < T < \delta$*, where* $\delta \in (0, A)$ *small enough. Then, there exist positive constants* C *and* s_0 *such that for all* $s \geq s_0$*, every solution* w *of* (2.9) *satisfies*

$$\int_Q (s\Theta k w_x^2 + s^3\Theta^3 \frac{(x-x_0)^2}{k} w^2) e^{2s\varphi} dtdadx \leq C \left(\int_q s^3\Theta^3 w^2 e^{2s\Phi} dtdadx + \int_0^1 \int_0^\delta w_T^2(a,x) dadx \right), \tag{2.50}$$

for all $s \geq s_0$ *and* δ *verifying* (2.4)*.*

Proof. Applying the inequality (2.31) to the function $h(t,a,x) = -\beta(t,a,x)w(t,0,x)$, we have the existence of two positive constants C and s_0 such that, for all $s \geq s_0$, the following inequality holds

$$\begin{aligned} &s^3 \int_Q \Theta^3 \frac{(x-x_0)^2}{k(x)} w^2 e^{2s\varphi} dtdadx + s \int_Q \Theta k(x) w_x^2 e^{2s\varphi} dtdadx \\ &\leq C \left(\int_Q \beta^2 w^2(t,0,x) e^{2s\Phi} dtdadx + \int_q s^3\Theta^3 w^2 e^{2s\Phi} dtdadx \right) \\ &\leq C \left(A\|\beta\|_\infty^2 \int_0^1 \int_0^T w^2(t,0,x) dtdx + \int_q s^3\Theta^3 w^2 e^{2s\Phi} dtdadx \right), \end{aligned} \tag{2.51}$$

using (2.4). On the other hand, integrating over the characteristics lines and after a careful calculus we obtain the following implicit formula of w solution of (2.5)

$$\begin{cases} w(t,a,\cdot) = \int_0^{A-a} S(A-a-l)\beta(t,A-l,\cdot)w(t,0,\cdot)dl, & \text{if } a > t + (A-T) \\ w(t,a,\cdot) = S(T-t)w_T(T+(a-t),\cdot) + \int_t^T S(l-t)\beta(l,a,\cdot)w(l,0,\cdot)dl, & \text{if } a \leq t + (A-T), \end{cases} \tag{2.52}$$

where $(S(t))_{t\geq 0}$ is the semi-group generated by the operator $A_2 w = (kw_x)_x - \mu w$.
Thus,

$$w(t,0,\cdot) = S(T-t)w_T(T-t,\cdot), \tag{2.53}$$

using the last hypothesis in (2.4) on β . Injecting this formula in (2.51) and using the fact $(S(t))_{t\geq 0}$ is a bounded semi-group, we get

$$s^3 \int_Q \Theta^3 \frac{(x-x_0)^2}{k(x)} w^2 e^{2s\varphi} dtdadx + s \int_Q \Theta k(x) w_x^2 e^{2s\varphi} dtdadx$$

$$\leq \widehat{C}\left(\int_0^1\int_0^T w_T^2(T-t,x)dtdx+\int_q s^3\Theta^3w^2e^{2s\Phi}dtdadx\right)$$
$$\leq \widehat{C}\left(\int_0^1\int_0^T w_T^2(m,x)dmdx+\int_q s^3\Theta^3w^2e^{2s\Phi}dtdadx\right)$$
$$\leq \widehat{C}\left(\int_0^1\int_0^\delta w_T^2(m,x)dmdx+\int_q s^3\Theta^3w^2e^{2s\Phi}dtdadx\right), \tag{2.54}$$

since $T\in(0,\delta)$ with $\delta\in(0,A)$ small enough and this achieves the proof of (2.50). □

3. **Observability inequality and null controllability results**

3.1. **Observability inequality result.** The objective of this paragraph is to reach the observability inequality of the adjoint system (2.5). To attain this purpose, we will combine the Carleman estimate (2.50) with the Hardy-Poincaré inequality stated in [22, Proposition 2.3] and arguing in a similar way as in [2]. Our observability inequality is given by the following proposition

Proposition 3.1. *Assume that the assumptions* (2.3), (2.4) *and* (2.7) *hold. Let* $A>0$ *and* $T>0$ *be given such that* $0<T<\delta$, *where* $\delta\in(0,A)$ *small enough. Then, there exists a positive constant* C_δ *such that for every solution* w *of* (2.5), *the following observability inequality holds*

$$\int_0^1\int_0^A w^2(0,a,x)dadx\leq C_\delta\left(\int_q w^2dtdadx+\int_0^1\int_0^\delta w_T^2(a,x)dadx\right). \tag{3.55}$$

Proof. Let w be a solution of (2.5). Then for $\kappa>0$ to be defined later, $\widetilde{w}=e^{\kappa t}w$ is a solution of

$$\begin{aligned}
&\frac{\partial\widetilde{w}}{\partial t}+\frac{\partial\widetilde{w}}{\partial a}+(k(x)\widetilde{w}_x)_x-(\mu(t,a,x)+\kappa)\widetilde{w}=-\beta\widetilde{w}(t,0,x),\\
&\widetilde{w}(t,a,1)=\widetilde{w}(t,a,0)=0,\\
&\widetilde{w}(T,a,x)=e^{\kappa T}w_T(a,x),\\
&\widetilde{w}(t,A,x)=0.
\end{aligned} \tag{3.56}$$

We point out that the parameter κ considered here is not the same as in (2.27). Multiplying the first equation of (3.56) by $\widetilde{w}$ and integrating by parts on $Q_t=(0,t)\times(0,A)\times(0,1)$. Then, one obtains

$$\begin{aligned}
&-\frac{1}{2}\int_{Q_A}\widetilde{w}^2(t,a,x)dadx+\frac{1}{2}\int_{Q_A}w^2(0,a,x)dadx+\frac{1}{2}\int_0^1\int_0^t\widetilde{w}^2(\tau,0,x)d\tau dx\\
&+\kappa\int_0^1\int_0^A\int_0^t\widetilde{w}^2(\tau,a,x)d\tau dadx\leq\frac{\|\beta\|_\infty^2}{4\epsilon'}\int_0^1\int_0^A\int_0^t\widetilde{w}^2(\tau,a,x)d\tau dadx\\
&+\epsilon' A\int_0^1\int_0^t\widetilde{w}^2(\tau,0,x)d\tau dx.
\end{aligned} \tag{3.57}$$

Thus, for $\kappa = \frac{\|\beta\|_\infty^2}{4\epsilon'}$ and $\epsilon' < \frac{1}{2A}$, one gets after integration over $(\frac{T}{4}, \frac{3T}{4})$

$$\int_{Q_A} w^2(0,a,x)dadx \leq C_{12}e^{2\kappa T}\int_{Q_A}\int_{\frac{T}{4}}^{\frac{3T}{4}} w^2(t,a,x)dadx. \tag{3.58}$$

On the other hand, let us prove that there exists a positive constant C_δ such that

$$\int_0^1\int_0^{\delta-\frac{3T}{4}}\int_{\frac{T}{4}}^{\frac{3T}{4}} w^2(t,a,x)dtdadx \leq C_\delta\int_0^1\int_0^\delta w_T^2(a,x)dadx. \tag{3.59}$$

For this purpose, we will use the implicit formula of w defined by (2.52) and we shall discuss the two cases, namely the case when $a > t + (A-T)$ and when $a \leq t + (A-T)$. In fact, if $a > t + (A-T)$ one has

$$\begin{aligned} w(t,a,\cdot) &= \int_0^{A-a} S(A-a-l)\beta(t,A-l,\cdot)w(t,0,\cdot)dl \\ &= \int_0^{A-a} S(A-a-l)\beta(t,A-l,\cdot)S(T-t)w_T(T-t,\cdot)dl, \end{aligned}$$

using (2.53). Since $(S(t))_{t\geq 0}$ is a bounded semi-group and $\beta \in L^\infty(Q)$, one can see that for $T \in (0,\delta)$

$$\begin{aligned} &\int_0^1\int_0^{\delta-\frac{3T}{4}}\int_{\frac{T}{4}}^{\frac{3T}{4}} w^2(t,a,x)dtdadx \\ &\leq \widetilde{C}_{10}\int_0^1\int_{\frac{T}{4}}^{\frac{3T}{4}} w_T^2(T-t,x)dtdx \\ &\leq \widetilde{C}_{10}\int_0^1\int_0^\delta w_T^2(m,x)dmdx, \end{aligned} \tag{3.60}$$

Now, if $a \leq t + (A-T)$ one has

$$\begin{aligned} w(t,a,\cdot) &= S(T-t)w_T(T+(a-t),\cdot) + \int_t^T S(l-t)\beta(l,a,\cdot)w(l,0,\cdot)dl \\ &= S(T-t)w_T(T+(a-t),\cdot) + \int_t^T S(l-t)\beta(l,a,\cdot)S(T-l)w_T(T-l,\cdot)dl. \end{aligned}$$

Thanks to the same argument employed to get (3.60), we conclude that

$$\begin{aligned} &\int_0^1\int_0^{\delta-\frac{3T}{4}}\int_{\frac{T}{4}}^{\frac{3T}{4}} w^2(t,a,x)dtdadx \\ &\leq 2\widetilde{C}_{11}\Big(\int_0^1\int_0^{\delta-\frac{3T}{4}}\int_{\frac{T}{4}}^{\frac{3T}{4}} w_T^2(T+(a-t),x)dtdadx \\ &+ \int_0^1\int_t^T w_T^2(T-l,x)dldx\Big). \end{aligned} \tag{3.61}$$

On one hand, we can check that

$$\int_0^1\int_0^{\delta-\frac{3T}{4}}\int_{\frac{T}{4}}^{\frac{3T}{4}} w_T^2(T+(a-t),x)dtdadx$$

$$\leq \int_0^1 \int_0^{\delta-\frac{3T}{4}} \int_{\frac{T}{4}}^{\frac{3T}{4}} w_T^2(a+m,x)dmdadx$$
$$\leq \int_0^1 \int_0^{\delta-\frac{3T}{4}} \int_{a+\frac{T}{4}}^{a+\frac{3T}{4}} w_T^2(z,x)dzdadx$$
$$\leq \int_0^1 \int_0^{\delta-\frac{3T}{4}} \int_0^{\delta} w_T^2(z,x)dzdadx$$
$$\leq \delta \int_0^1 \int_0^{\delta} w_T^2(z,x)dzdx. \tag{3.62}$$

On the other hand, we have the following inequality

$$\int_0^1 \int_t^T w_T^2(T-l,x)dldx = \int_0^1 \int_0^{T-t} w_T^2(z,x)dzdx$$
$$\leq \int_0^1 \int_0^{\delta} w_T^2(z,x)dzdx. \tag{3.63}$$

Combining the inequalities (3.61), (3.62) and (3.63) we get

$$\int_0^1 \int_0^{\delta-\frac{3T}{4}} \int_{\frac{T}{4}}^{\frac{3T}{4}} w^2(t,a,x)dtdadx$$
$$\leq \widetilde{C}_{12} \int_0^1 \int_0^{\delta} w_T^2(z,x)dzdx.$$

Subsequently, (3.59) occurs in both studied cases. Therefore, in the light of inequality (3.58) we conclude that

$$\int_{Q_A} w^2(0,a,x)dadx \leq \widetilde{C}_{13} \int_0^1 \int_0^{\delta} w_T^2(a,x)dadx$$
$$+\frac{2e^{2\kappa T}}{T} \int_0^1 \int_{\delta-\frac{3T}{4}}^{A} \int_{\frac{T}{4}}^{\frac{3T}{4}} w^2(t,a,x)dtdadx. \tag{3.64}$$

Now, let p defined by (2.18). Then, using the hypotheses (2.3) on k the function $x \mapsto \frac{(x-x_0)^2}{p(x)}$ is nonincreasing in the left of x_0 and nondecreasing in the right of x_0. Hence, applying Hardy-Poincaré inequality (see [22, Proposition 2.3]) and taking into account the definition of φ stated in (2.6) we have

$$\int_{Q_A} w^2(0,a,x)dadx \leq \widetilde{C}_{13} \int_0^1 \int_0^{\delta} w_T^2(a,x)dadx$$
$$+C_\delta^{13} \int_0^1 \int_{\delta-\frac{3T}{4}}^{A} \int_{\frac{T}{4}}^{\frac{3T}{4}} s\Theta k(x) w_x^2(t,a,x) e^{2s\varphi} dtdadx. \tag{3.65}$$

Therefore, using Carleman estimate (2.50) we infer

$$\int_{Q_A} w^2(0,a,x)dadx \leq \widetilde{C}_\delta^{15} \left(\int_q s^3\Theta^3 w^2 e^{2s\Phi} dtdadx + \int_0^1 \int_0^{\delta} w_T^2(a,x)dadx \right),$$

and then the proof is finished using the fact that $\sup_{(t,a,x)\in Q} s^d\Theta^d e^{2s\Phi} < +\infty \ , \forall d \in \mathbb{R}$. □

3.2. **Null controllability result.** In the previous paragraph, we obtained the observability inequality of system (2.5). Such a tool will be very useful to prove the null controllability of the model (1.1) in the case where $T \in (0, \delta)$ as we emphasized in the introduction. Our main result is provided in the following theorem

Theorem 3.2. *Assume that the dispersion coefficient k satisfies* (2.3) *and the natural rates β and μ verify* (2.4). *Let $A, T > 0$ be given such that $0 < T < \delta$, where $\delta \in (0, A)$ small enough. For all $y_0 \in L^2(Q_A)$, there exists a control $\vartheta \in L^2(q)$ such that the associated solution of* (1.1) *verifies*

$$y(T, a, x) = 0, \qquad a.e.\ in\ (\delta, A) \times (0, 1). \tag{3.66}$$

Furthermore, there exists a positive constant C_{10} which depends on δ such that ϑ satisfies the following inequality.

$$\int_q \vartheta^2(t, a, x) dt da dx \leq C_{10} \int_{Q_A} y_0^2(a, x) da dx. \tag{3.67}$$

C_{10} is called the control cost.

Before proceeding to the proof of Theorem 3.2, we shall make the following remark:

Remark 3.3. *The inequality* (3.67) *shows us clearly that the control that we are looking for depends on δ and the initial distribution y_0.*

Proof. Let $\varepsilon > 0$ and consider the following cost function

$$J_\varepsilon(\vartheta) = \frac{1}{2\varepsilon} \int_0^1 \int_\delta^A y^2(T, a, x) da dx + \frac{1}{2} \int_q \vartheta^2(t, a, x) dt da dx.$$

We can prove that J_ε is continuous, convex and coercive. Then, it admits at least one minimizer ϑ_ε and we have

$$\vartheta_\varepsilon = -w_\varepsilon(t, a, x)\chi_\omega(x) \quad \text{in } Q, \tag{3.68}$$

with w_ε is the solution of the following system

$$\begin{aligned}
\frac{\partial w_\varepsilon}{\partial t} + \frac{\partial w_\varepsilon}{\partial a} + (k(x)(w_\varepsilon)_x)_x - \mu(t, a, x) w_\varepsilon &= -\beta w_\varepsilon(t, 0, x) && \text{in } Q, \\
w_\varepsilon(t, a, 1) = w_\varepsilon(t, a, 0) &= 0 && \text{on } (0, T) \times (0, A), \\
w_\varepsilon(T, a, x) &= \frac{1}{\varepsilon} y_\varepsilon(T, a, x)\chi_{(\delta, A)}(a) && \text{in } Q_A, \\
w_\varepsilon(t, A, x) &= 0 && \text{in } Q_T,
\end{aligned} \tag{3.69}$$

and y_ε is the solution of the system (1.1) associated to the control ϑ_ε. Multiplying (3.69) by y_ε, integrating over Q, using (3.68) and the Young inequality we obtain

$$\begin{aligned}
&\frac{1}{\varepsilon} \int_0^1 \int_\delta^A y_\varepsilon^2(T, a, x) da dx + \int_q \vartheta_\varepsilon^2(t, a, x) dt da dx \\
= &\int_{Q_A} y_0(a, x) w_\varepsilon(0, a, x) da dx \\
&\leq \frac{1}{4C_\delta} \int_{Q_A} w_\varepsilon^2(0, a, x) da dx + C_\delta \int_{Q_A} y_0^2(a, x) da dx,
\end{aligned}$$

with C_δ is the constant of the observability inequality (3.55). This again leads to

$$\frac{1}{\varepsilon}\int_0^1\int_\delta^A y_\varepsilon^2(T,a,x)dadx+\int_q \vartheta_\varepsilon^2(t,a,x)dtdadx\leq \frac{1}{4}\int_q w^2dtdadx+C_\delta\int_{Q_A} y_0^2(a,x)dadx.$$

Keeping in the mind (3.68), we conclude that

$$\frac{1}{\varepsilon}\int_0^1\int_\delta^A y_\varepsilon^2(T,a,x)dadx+\frac{3}{4}\int_q \vartheta_\varepsilon^2(t,a,x)dtdadx\leq C_\delta\int_{Q_A} y_0^2(a,x)dadx. \tag{3.70}$$

Hence, it follows that

$$\begin{cases}\int_0^1\int_\delta^A y_\varepsilon^2(T,a,x)dadx\leq \varepsilon C_\delta\int_{Q_A} y_0^2(a,x)dadx\\ \int_q \vartheta_\varepsilon^2(t,a,x)dtdadx\leq \frac{4C_\delta}{3}\int_{Q_A} y_0^2(a,x)dadx.\end{cases} \tag{3.71}$$

Then, we can extract two subsequences of y_ε and ϑ_ε denoted also by ϑ_ε and y_ε that converge weakly towards ϑ and y in $L^2(q)$ and $L^2((0,T)\times(0,A);H_k^1(0,1))$ respectively. Now, by a variational technic, we prove that y is a solution of (1.1) corresponding to the control ϑ and, by the first estimate of (3.71), y satisfies (3.66) for $T\in(0,\delta)$ and this shows our claimed Theorem3.2 □

4. **Appendix**

As we said in the introduction, this Appendix is concerned with a result which plays an important role to show the ω-Carleman estimate associated to the full adjoint system (2.5) namely the Caccioppoli's inequality which is stated in the following lemma

Lemma 4.1. *Let $\omega'\subset\subset\omega$ and w be the solution of (2.9). Suppose that $x_0\notin\overline{\omega'}$. Then, there exists a positive constant C such that w verifies*

$$\int_{\omega'}\int_0^A\int_0^T w_x^2e^{2s\varphi}dtdadx\leq C\left(\int_q s^2\Theta^2w^2e^{2s\varphi}dtdadx+\int_q h^2e^{2s\varphi}dtdadx\right). \tag{4.72}$$

Proof. Define the following smooth cut-off function $\zeta:\mathbb{R}\to\mathbb{R}$

$$\begin{cases}0\leq\zeta(x)\leq 1, & x\in\mathbb{R},\\ \zeta(x)=0, & x<x_1 \text{ and } x>x_2,\\ \zeta(x)=1, & x\in\omega'.\end{cases} \tag{4.73}$$

For the solution w of (2.9), we have

$$\begin{aligned}0&=\int_0^T\frac{d}{dt}\left[\int_0^1\int_0^A\zeta^2e^{2s\varphi}w^2dadx\right]dt\\ &=2s\int_0^1\int_0^A\int_0^T\zeta^2\varphi_tw^2e^{2s\varphi}dtdadx+2\int_0^1\int_0^A\int_0^T\zeta^2ww_te^{2s\varphi}dtdadx\\ &=2s\int_0^1\int_0^A\int_0^T\zeta^2\varphi_tw^2e^{2s\varphi}dtdadx+2\int_0^1\int_0^A\int_0^T\zeta^2w(-(kw_x)_x-w_a+h+\mu w)e^{2s\varphi}dtdadx.\end{aligned}$$

Then, integrating by parts we obtain

$$2\int_Q k\zeta^2e^{2s\varphi}w_x^2dtdadx=-2s\int_Q\zeta^2w^2\psi(\Theta_a+\Theta_t)e^{2s\varphi}dtdadx-2\int_Q\zeta^2whe^{2s\varphi}dtdadx$$

$$-2\int_Q \zeta^2\mu w^2 e^{2s\varphi}dtdadx + \int_Q (k(\zeta^2 e^{2s\varphi})_x)_x w^2 dtdadx.$$

On the other hand, by the definitions of ζ, $\underline{\psi}$ and Θ, thanks to Young inequality, taking s quite large and using the fact that $x_0 \notin \overline{\omega'}$, one can prove the existence of a positive constant c such that

$$2\int_Q k\zeta^2 e^{2s\varphi} w_x^2 dtdadx \geq 2\min_{x\in\omega'} k(x)\int_{\omega'}\int_0^A\int_0^T w_x^2 e^{2s\varphi}dtdadx,$$

$$\int_Q (k(\zeta^2 e^{2s\varphi})_x)_x w^2 dtdadx \leq c\int_\omega\int_0^A\int_0^T s^2\Theta^2 w^2 e^{2s\varphi}dtdadx,$$

$$-2s\int_Q \zeta^2 w^2\psi(\Theta_a+\Theta_t)e^{2s\varphi}dtdadx \leq c\int_\omega\int_0^A\int_0^T s^2\Theta^2 w^2 e^{2s\varphi}dtdadx,$$

$$-2\int_Q \zeta^2 whe^{2s\varphi}dtdadx \leq c\left(\int_\omega\int_0^A\int_0^T s^2\Theta^2 w^2 e^{2s\varphi}dtdadx + \int_\omega\int_0^A\int_0^T h^2 e^{2s\varphi}dtdadx\right),$$

$$-2\int_Q \zeta^2\mu w^2 e^{2s\varphi}dtdadx \leq c\int_\omega\int_0^A\int_0^T s^2\Theta^2 w^2 e^{2s\varphi}dtdadx.$$

This all together imply that there is $C>0$ such that

$$\int_{\omega'}\int_0^A\int_0^T w_x^2 e^{2s\varphi}dtdadx \leq C\left(\int_q s^2\Theta^2 w^2 e^{2s\varphi}dtdadx + \int_q h^2 e^{2s\varphi}dtdadx\right).$$

Thus, the proof is achieved. □

Remark 4.2. *The Lemma 4.1 remains true for any function* $\pi \in C([0,1],(-\infty,0)) \cap C^1([0,1]\backslash\{x_0\},(-\infty,0))$ *and verifying*

$$|\pi_x| \leq \frac{c}{\sqrt{k}}, \qquad \text{for } x\in[0,1]\backslash\{x_0\}, \tag{4.74}$$

where $c>0$. *see* [22, Proposition 4.2] *for more details.*

Acknowledgements
The authors would like to thank deeply the anonymous referee and the Professors B. Ainseba and L. Maniar for their fruitful and several remarks which allow us to realize this work.

References

[1] B. Ainseba, Corrigendum to "Exact and approximate controllability of the age and space population dynamics structured model [J. Math. Anal. Appl. 275 (2002), 562-574]", J. Math. Anal. Appl. 393 (2012), 328.
[2] B. Ainseba, Exact and approximate controllability of the age and space population dynamics structured model, J. Math. Anal. Appl. 275 (2002), 562-574.
[3] B. Ainseba and S. Anita, Internal exact controllability of the linear population dynamics with diffusion, Electronic Journal of Differential Equations, 2004(2004), 1-11.
[4] B. Ainseba and S. Anita, Local exact controllability of the age-dependent population dynamics with diffusion, Abstr. Appl. Anal. 6 (2001), 357-368.
[5] B. Ainseba, Y. Echarroudi and L. Maniar, Null controllability of a population dynamics with degenerate diffusion, Journal of Differential and Integral Equations, 26 (2013), 1397-1410.

[6] F. Alabau-Boussouira, P. Cannarsa and G. Fragnelli, Carleman estimates for degenerate parabolic operators with applications to null controllability, J. evol.equ. 6 (2006), 161-204.
[7] V. Barbu, M. Iannelli and M. Martcheva, On the controllability of the Lotka-McKendrick model of population dynamics, J. Math. Anal. Appl. 253 (2001), 142-165.
[8] A. Bátkai, P. Csomós, B. Farkas and G. Nickel, Operator splitting for non-autonomous evolution equations, Journal of Functional Analysis 260 (2011) 21632190.
[9] A. Bátkai and S. Piazzera, Semigroups for delay equations, Research Notes in Mathematics ; 10, Wellesley, MA : Peters, 2005. - 259 p.
[10] I. Boutaayamou, G. Fragnelli and L. Maniar, Carleman estimates for parabolic equations with interior degeneracy and Neumann boundary conditions, Accepted for publication in J. Anal. Math.
[11] P. Cannarsa, P. Martinez and J. Vancostenoble, *Null controllability of degenerate heat equations,* Adv. Differential Equations. 10(2005), pp. 153-190.
[12] P. Cannarsa, G. Fragnelli, Null controllability of semilinear degenerate parabolic equations in bounded domains, Electron. J. Differential Equations 2006 (2006), 120.
[13] P. Cannarsa, G. Fragnelli, D. Rocchetti, Controllability results for a class of one- dimensional degenerate parabolic problems in nondivergence form, J. Evol. Equ. 8 (2008), 583616.
[14] P. Cannarsa, G. Fragnelli, D. Rocchetti, Null controllability of degenerate parabolic operators with drift, Netw. Heterog. Media 2 (2007), 693713.
[15] P. Cannarsa, G. Fragnelli, J. Vancostenoble, Regional controllability of semilinear degenerate parabolic equations in bounded domains, J. Math. Anal. Appl. 320 (2006), 804818.
[16] P. Cannarsa, G. Fragnelli, J. Vancostenoble, Linear degenerate parabolic equations in bounded domains: controllability and observability, IFIP Int. Fed. Inf. Process. 202 (2006), 163173, Springer, New York.
[17] Y. Echarroudi and L. Maniar, Null controllability of a model in population dynamics, Electronic Journal of Differential Equations, 2014 (2014), No. 240, 1-20.
[18] G. Fragnelli, An age dependent population equation with diffusion and delayed birth process, International Journal of Mathematics and Mathematical Sciences, 20 (2005), 3273-3289.
[19] G. Fragnelli, Null controllability of degenerate parabolic equations in non divergence form via Carleman estimates, Discrete Contin. Dyn. Syst. Ser. S 6 (2013), 687701.
[20] G. Fragnelli, A. Idrissi and L. Maniar, The asymptotic behaviour of a population equation with diffusion and delayed birth process, Discrete and Continuous Dynamical Systems-Series B, 7 (2007), No. 4, 735-754.
[21] G. Fragnelli, P. Martinez and J. Vancostenoble, Qualitative properties of a population dynamics describing pregnancy, Math. Models Methods Appl. Sci. 15, 507 (2005). DOI: 10.1142/S0218202505000455
[22] G. Fragnelli and D. Mugnai, Carleman estimates and observability inequalities for parabolic equations with interior degeneracy, Advances in Nonlinear Analysis 08/2013; 2(4):339378. DOI: 10.1515/anona-2013-0015.
[23] G. Fragnelli and D. Mugnai, Carleman estimates, observability inequalities and null controllability for interior degenerate non smooth parabolic equations, to appear in Mem. Amer. Math. Soc. ArXiv: 1508.04014.
[24] G. Fragnelli, G. Ruiz Goldstein, J.A. Goldstein, S. Romanelli, Generators with interior degeneracy on spaces of L2 type, Electron. J. Differential Equations 2012 (2012), 1-30.
[25] G. Fragnelli and L. Tonetto, A population equation with diffusion, J. Math. Anal. Appl., 289 (2004), 90-99.
[26] A. V. Fursikov and O. Yu. Imanuvilov, Controllability of Evolution Equations, Lecture Notes Series, vol. 34, Seoul National University Research Institute of Mathematics Global Analysis Research Center, Seoul, 1996.
[27] M. Langlais, A nonlinear problem in age-dependent population diffusion, Siam J. Math. Anal. 16 (1985), 510-529.
[28] S. Piazzera, An age-dependent population equation with delayed birth process, Mathematical Methods in the Applied Sciences, Vol. 27, Issue 4, pages 427439, 10 March 2004.
[29] M. A. Pozio and A. Tesei, Degenerate parabolic Problems in population dynamics, Japan Journal of Applied Mathematics, December 1985, 2:351

[30] G. Nickel, Evolution semigroups for nonautonomous Cauchy problems, Abstr. Appl. Anal. 2 (1997) 7395.
[31] A. Rhandi and R. Schnaubelt, Asymptotic behaviour of a non-autonomous population equation with diffusion in L^1, Discrete Contin. Dynam. Systems 5 (1999), 663-683.
[32] G. F. Webb, Population models structured by age, size, and spatial position. Structured population models in biology and epidemiology, 149, Lecture Notes in Math. 1936, Springer, Berlin, 2008.